

Structural Analysis and Characteristics of Oligosaccharides Isolated from Fermented Beverage of Plant Extract

Hideki Okada^{1*} • Naoki Kawazoe¹ • Akira Yamamori¹ • Shuichi Onodera² • Norio Shiomi²

¹ Ohtakakohso, Co., Ltd., 1-22-10, Sakura, Otaru 047-0193, Japan

² Department of Food and Nutrition Sciences, Graduate School of Dairy Science Research, Rakuno Gakuen University, 582, Bunkyoudai-Midorimachi, Ebetsu 069-8501, Japan

Corresponding author: * oelab@ohtakakohso.co.jp

ABSTRACT

A fermented beverage of plant extract was prepared from about 50 kinds of fruits and vegetables. Natural fermentation was conducted by lactic acid bacteria (*Leuconostoc* spp.) and yeast (*Zygosaccharomyces* spp. and *Pichia* spp.). Eighteen kinds of oligosaccharides were isolated from this beverage, and their structures were confirmed by methylation analysis, MALDI-TOF-MS and NMR measurements. In these saccharides, eight novel oligosaccharides were found to be constructed by di- and trisaccharides with the fructosyl residue of pyranose form, and other trisaccharides with fructosyl residues of sucrose bond with the β -D-galactose and β -D-glucose. The characteristics of one of novel saccharide, *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose (Fp2-6G) were investigated, and it was shown to be non-cariogenicity and have low digestibility. Furthermore, the saccharide was selectively used by beneficial bacteria, *Bifidobacterium adolescentis* and *B. longum*, but was not used by unfavorable bacteria, *Clostridium perfringens*, *Escherichia coli* and *Enterococcus faecalis* that produce mutagenic substances.

Keywords: β -D-fructopyranoside, natural fructopyranoside, novel oligosaccharide

Abbreviations: Fp2-6G, *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose

CONTENTS

INTRODUCTION.....	1
METHODOLOGY.....	2
Preparation of fermented beverage of plant extract.....	2
High performance anion-exchange chromatography (HPAEC).....	2
<i>p</i> -Aminobenzoic acid ethyl ester conversion method.....	2
Methylation and methanolysis.....	2
Gas liquid chromatography.....	2
GC-MS analysis.....	2
MALDI-TOF-MS.....	2
NMR measurement.....	2
Cariogenicity of <i>O</i> - β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose (Fp2-6G).....	3
RESEARCH FINDINGS.....	3
Isolation of saccharides from fermented beverage of plant extract.....	3
Structure analysis of saccharides.....	3
Characteristics of saccharide 2, <i>O</i> - β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose (Fp2-6G).....	6
SUPPLEMENTATION.....	6
ACKNOWLEDGEMENTS.....	6
REFERENCES.....	6

INTRODUCTION

The extract from 50 kinds of fruits and vegetables was fermented to produce a new beverage (Okada *et al.* 2005, 2006). The juices were extracted using sucrose-osmotic pressure in a cedar barrel for one week and were fermented by lactic acid bacteria (*Leuconostoc* spp.) and yeast (*Zygosaccharomyces* spp. and *Pichia* spp.). The fermented beverage showed scavenging activity against 1,1'-diphenyl-2-picrylhydrazyl (DPPH) radical, and reduced significantly the ethanol-induced damage of gastric mucosa in rat (Okada *et al.* 2005). Analysis by high performance anion-exchange chromatography (HPAEC) showed that this beverage contained high levels of saccharides, estimated between 550 and 590 gL⁻¹, mainly glucose and fructose, and a small

amount of undetermined oligosaccharides. Recently, it was reported that different positions of the glycosidic linkage of oligosaccharide isomers affect the isomer's physiological and physical properties (Kohmoto *et al.* 1988; Murosaki *et al.* 1999, 2002). The development of HPLC analyses with high sensitivity and separation ability enable the detection and isolation of oligosaccharides in fermented beverages. We already reported that 18 kinds of oligosaccharides were isolated from this beverage, in these saccharides, eight novel oligosaccharides [saccharide 2 (Okada *et al.* 2006), 10, 14 (Kawazoe *et al.* 2008a), 12, 13 (Kawazoe *et al.* 2008b), 8, 9, 16 (Okada *et al.* 2009)] were found.

In this paper, we report on the isolation and structural confirmation of the saccharides from the fermented beverage of plant extracts.

METHODOLOGY

Preparation of fermented beverage of plant extract

For preparation of initial juice, 50 kinds of fruits and vegetables were used to produce the final extract as detailed in Okada *et al.* (2005). Fruits and vegetables, obtained from local markets of Hokkaido region, were sorted for the absence of defects, washed and immediately processed for juice extraction. The fruits and vegetables were cut, sliced or diced into small pieces, mixed and placed in cedar barrels. Afterwards, the equivalent weight of sucrose was added to samples, mixed well to allow high samples-sucrose contact, and then barrels were left at room temperature for one week. The juice exudates were then separated – without compression – from solids and used for fermentation. Almost all sucrose was hydrolyzed to glucose and fructose during fermentation (glucose 30%, fructose 25%). The fermented beverage was obtained by incubation of the juice at 37°C in the dark by natural fermentation using yeast (*Zygosaccharomyces* spp. and *Pichia* spp.) and lactic acid bacteria (*Leuconostoc* spp.). After 7 days, the fermented beverage was kept in a closed enameled tank (5,000 L) at 37°C for 6-10 months for additional maturation and aging to finally obtain a brown and slightly sticky liquid (Super Ohtaka, Ohtakakohso, Co., Ltd. Hokkaido).

The pH of plant extract reached about 3.2 after 7 days fermentation (Fig. 1).

High performance anion-exchange chromatography (HPAEC)

The oligosaccharides were analyzed using a Dionex Bio LC Series apparatus (Sunnyvale, CA) equipped with an HPLC carbohydrate column (Carbo Pac PA1, inert styrene divinyl benzene polymer, Sunnyvale, CA) and a pulsed amperometric detector (PAD: Rocklin and Pohl 1983; Johnson 1986). The gradient was established by mixing eluant A (150 mM NaOH) with eluant B (500 mM sodium acetate in 150 mM NaOH) as follows: 0-1 min, 25 mM; 1-2 min, 25-50 mM; 2-20 min, 50-200 mM; 20-22 min, 500 mM; 22-30 min, 25 mM; using a flow rate through the column of 1.0 mL/min. The applied PAD potentials for E1 (500 ms), E2 (100 ms) and E3 (50 ms) were 0.1, 0.6 and -0.6V, respectively, and the output range was 1 μ C.

p-Aminobenzoic acid ethyl ester conversion method

Conversion of the saccharides at the reducing end with p-Aminobenzoic acid ethyl ester (ABEE: Seikagaku Biobusiness Co., Tokyo) was carried out according to a prior method (Yasuno *et al.* 1997; Kobayashi *et al.* 2002). Ten μ L of standard saccharide solution was added to an ABEE reagent solution (40 μ L). The mixture was incubated at 80°C for 1 h. Distilled water (0.2 mL) and chloroform (0.2 mL) were added and mixture was centrifuged at 2,000 \times g for 1 min, aqueous layer was diluted (100-fold) with water and subjected to HPLC analysis. ABEE-converted saccharide was detected by UV at 305 nm.

Methylation and methanolysis

Methylation of the oligosaccharides was carried out by the method of Hakomori (Hakomori 1964).

The permethylated saccharides were methanolized by heating with 1.5% methanolic hydrochloric acid at 96°C for 10 or 180 min. The reaction mixture was treated with Amberlite IRA-410 (OH⁻: Organo Co., Tokyo) to remove hydrochloric acid, and evaporated *in vacuo* to dryness. The resulting methanolysate was dissolved in a small volume of methanol and analyzed using gas liquid chromatography (GLC).




Fig. 1 Fermentation process of plant extracts. Symbols: \circ , pH; \blacksquare , colony-forming units/g.

Gas liquid chromatography

For the analysis of the methanolysate, GLC was carried out using a Shimadzu GC-8A gas chromatograph equipped with a glass column (2.6 mm \times 2 m) packed with 15% butane 1,4-diol succinate polyester on acid-washed Celite at 175°C. The flow rate of the nitrogen gas carrier was 40 mL/min.

GC-MS analysis

GC-MS analysis (Ciucanu and Costello 2003) was performed using JMS-AX500 mass spectrometer (JEOL, Japan) using a DB-17HT capillary column (30 m \times 0.25 mm I.D., J&W Scientific, USA). Injection temperature was 200°C. The column temperature was kept at 50°C for 2 min after sample injection, increased to 150°C at 50°C/min, kept at 150°C for 1 min, then increased to 250°C at 4°C/min. The mass spectra were recorded in the electron ionization (EI) mode.

MALDI-TOF-MS

MALDI-TOF-MS spectra were measured using a Shimadzu-Kratos mass spectrometer (KOMPACT Probe) in positive ion mode with 2, 5-dihydroxybenzoic acid as a matrix. Ions were formed by a pulsed UV laser beam (nitrogen laser, 337 nm). Calibration was done using 1-kestose as an external standard. 1-Kestose was prepared from sucrose using a *Scopulariopsis brevicaulis* enzyme (Takeda *et al.* 1994).

NMR measurement

The saccharides were dissolved in 0.4 mL D₂O. NMR spectra were recorded at 27°C with a Bruker AMX 500 spectrometer (¹H 500 MHz, ¹³C 125 MHz) equipped with a 5 mm diameter C/H dual (1D spectra) and TXI probe (2D spectra). Chemical shifts of ¹H (δ_H) and ¹³C (δ_C) in ppm were determined relative to the external standard of sodium [2, 2, 3, 3-²H₄]-3-(trimethylsilyl)-propanoate in D₂O (δ_H 0.00 ppm) and 1,4-dioxane (δ_C 67.40 ppm) in D₂O, respectively. ¹H-¹H COSY (Aue *et al.* 1975; von Kienlin *et al.* 1991), HSQC (Willker *et al.* 1993), HMBC (Bax and Summers 1986; Hurd and John 1991), and CT-HMBC (Bax and Summers 1986; Hurd and John 1991) spectra were obtained using gradient selected pulse sequences. The phase-sensitive HSQC-TOCSY (Domke 1991; Willker *et al.* 1993) and CH₂-selected E-HSQC-TOCSY (Yamamori *et al.* 2002) spectra were determined with the sequence including inversion of direct resonance (IDR). The TOCSY mixing time (108 ms) was composed of MLEV-17 composite pulses guarded by a trim pulse (2.5 ms). The coupling patterns of overlapped ¹H were analyzed by SPT method (Pachler and Wessels 1973; Uzawa and Yoshida 2004).

Cariogenicity of *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose (Fp2-6G)

The cariogenicity was performed according to the method of Miyamura (1973). The oral cavity of a subject was rinsed with tap water and then the subject gargled with distilled water. The saliva naturally secreted was collected and well shaken at room temperature. A 0.5 mL aliquot of 1% Fp2-6G, palatinose, glucose or distilled water was then added to a mixture of 1.5 mL of this fresh saliva and 0.5 mL of a brain heart infusion broth (Eiken Chemical Co., Ltd. Tokyo). These mixtures were incubated at 37°C to show the time-dependent pH variation of the cariogenicity. The same experiment was conducted with *Streptococcus mutans* JCM 5705 instead of saliva (the same experiment was repeated three times).

RESEARCH FINDINGS

Isolation of saccharides from fermented beverage of plant extract

The fermented beverage of plant extract (5 kg) was loaded onto to a carbon-Celite [1: 1; Charcoal (Wako Pure Chemical Industries, Ltd., Osaka, Japan) and Celite-535 (Nakalai Chemical Industries, Ltd., Osaka, Japan)] column (4.5 \times 35 cm) and successively eluted with water (14 L), 5% ethanol (30 L) and 30% ethanol. Almost all of glucose and fructose were eluted with water (4L) and then saccharides 1 and 2 were eluted with water (5-6L, Fig. 2a). Fractions including saccharides 1 and 2 were concentrated *in vacuo* and freeze-dried to give 1.25 g. Finally, the yield and purity of saccharide 2 were 49 and 98%, respectively. Saccharides 3, 4 and 5 were eluted with 1 L of 5% ethanol (Fig. 2b). Saccharide 6 was obtained by additional elution (1L) of 5% ethanol (Fig. 2c). Both fractions containing saccharides 5 and 6 were also concentrated *in vacuo* and freeze-dried to give 0.97 and 1.70 g, respectively. Saccharides 7-18 were eluted with 30% ethanol (1-2L, Fig. 2d, 0.84 g). Subsequently, those fractions were successfully purified repeatedly using an HPLC system (Tosoh, Tokyo, Japan) equipped with an Amide-80 column (7.8 mm \times 30 cm, Tosoh, Tokyo, Japan) at 80°C, and eluted with 80% acetonitrile at 2.0 mL/min, and using refractive index detection. Furthermore, those saccharides were purified using a HPLC with an ODS-80Ts column (4.6 mm \times 25 cm, Tosoh, Tokyo, Japan) at room temperature.

Structure analysis of saccharides

The saccharides isolated from the fermented beverage were identified using high performance anion-exchange chromatography (HPAEC: Shiomi *et al.* 1991; Okada *et al.* 2003), MALDI-TOF-MS and NMR.




Fig. 2 Isolation of saccharides from plant extract by Amide-80 column chromatography. Fermented beverage of plant extract was loaded onto to a carbon-Celite column and successively eluted with water (A), 5% ethanol (B, C) and 30% ethanol (D).

Saccharides 1, 3, 4, 5, 6, 7, 11, 15, 17 and 18 corresponded to authentic saccharides, *O*- β -D-fructofuranosyl-(2 \rightarrow 6)-glucose, maltose, sophorose, gentiobiose, laminaribiose, maltotriose, raffinose, panose, *O*- β -D-glucopyranosyl-(1 \rightarrow 6)-*O*- β -D-glucopyranosyl-(1 \rightarrow 3)-D-glucopyranose and gentiotriose, respectively. Sophorose, gentiobiose and laminaribiose were already known to be synthesized enzymatically under highly concentrated solution of glucose and the presence of yeast and *Penicillium* β -glucosidase (Villa and Phaff 1979; Fujimoto *et al.* 1988; Rastall *et al.* 1992). *O*- β -D-Glucopyranosyl-(1 \rightarrow 6)-*O*- β -D-glucopyranosyl-(1 \rightarrow 3)-D-glucopyranose was also reported to be synthesized by β -glucosidase (Kawamura *et al.* 1971). The saccharides isolated from the fermented beverage of plant extracts were not detected before fermentation. Therefore, these saccharides were estimated to be synthesized by action of yeast β -glucosidase during fermentation. Although fructofuranosyl-(2 \rightarrow 6)-glucose was obtained from partial hydrolysate of neokestose, it was presumed to be produced by reverse-reaction of invertase of yeast.

The $t_{R,sucrose}$ values of HPAEC of saccharides 2, 8, 9, 10, 12, 13, 14 and 16 did not correspond to those of any authentic saccharides. The degree of polymerization (DP) of

Table 1 Gas-liquid chromatographic analysis of methanolysates of permethylated saccharides.

Methanolysate origin	Relative retention time ^{a)}								
Saccharide 2	0.99			1.46		2.44		3.50	
Saccharide 8		1.00	1.07	1.30	1.48	2.59		3.50	
Saccharide 9			1.07	1.29		2.58	3.22	3.59	4.73
Saccharide 10	0.97			1.43		2.51	3.26	3.63	4.77
Saccharide 12		1.01			1.41	2.64			3.94
Saccharide 13		1.01			1.42	1.74	2.64		3.97
Saccharide 14	0.92	1.04		1.46	1.46				
Saccharide 16		0.97			1.47				
Fructose	0.93		1.07	1.30	1.42				
1-Kestose			1.08	1.28	1.49		2.65		3.96
Raffinose			1.04	1.34		1.69	2.63	3.74	
Laminaribiose	0.98				1.49			3.31	4.88
Methyl-2,3,4,6-tetra- <i>O</i> -methyl- β -D-glucopyranoside	1.00								
Methyl-2,3,4,6-tetra- <i>O</i> -methyl- β -D-galactopyranoside						1.76			

a) Retention time of methyl 2,3,4,6-tetra-*O*-methyl- β -D-glucopyranoside=1.

saccharide 2 was established as 2 by measurements of $[M+Na]^+$ ions (m/z : 365) using TOF-MS and similar analyses of saccharides 8, 9, 10, 12, 13, 14 and 16 established all their DP as 3 (m/z : 527). From the GC analysis, relative retention times of the methanolysate of the permethylated saccharide were investigated [t_R (relative retention time; retention time of methyl 2, 3, 4, 6-tetra-*O*-methyl- β -D-glucopyranoside = 1.0; retention time, 9.60 min)]. The methanolysate of permethylated saccharide 2 exhibited four peaks (Table 1), two of which corresponded to methyl 2,3,4-tri-*O*-methyl-D-glucopyranoside (t_R , 2.44 and 3.50). From this finding, saccharide 2 was presumed to be fructosyl (2 \rightarrow 6) glucose. However, other peaks (t_R , 0.99 and 1.46) did not correspond to methyl 1,3,4,6-tetra-*O*-methyl-D-fructofuranoside (t_R , 1.04 and 1.34) from permethylated raffinose. Because the peaks were estimated to correspond to methyl 1,3,4,5-tetra-*O*-methyl-D-fructopyranoside, we tried to prepare methyl fructopyranoside from-D-fructose. The permethylated fructose and methanolysate of permethylated saccharide 2 were analyzed by GC-MS. The permethylated fructose gave four peaks, that were confirmed to be methyl 1,3,4,6-tetra-*O*-methyl- β - and α -D-fructofuranoside (retention time, 8.34 and 8.56 min) and methyl 1,3,4,5-tetra-*O*-methyl- β - and α -D-fructopyranoside (9.14 and 10.57 min), respectively from the retention time and pattern of fragmentation (Ciucanu and Costello 2003). The methanolysate from permethylated saccharide 2 exhibited two peaks (9.14 and 10.63 min) corresponding to the α/β isomers methyl 1,3,4,5-tetra-*O*-methyl-D-fructopyranoside. No peaks corresponding to methyl 1,3,4,5-tetra-*O*-methyl-D-fructopyranoside were detected from the methanolysate of permethylated raffinose. From these findings as above, saccharide 2 was proved to be D-fructopyranosyl-(2 \rightarrow 6)-D-glucose.

Subsequently, NMR analysis was done. The HSQC-TOCSY spectrum revealed the 1H and ^{13}C signals of each β -Glc, α -Glc, and Fru residues. The COSY spectrum assigned the spin systems of these residues; from H-1 to H-6 in β -Glc, from H-1 to H-5 of α -Glc, and from H-4 to H-6 in Fru. The corresponding ^{13}C signals were assigned by HSQC spectrum. The assignment of the remaining signals, the position of the glycosidic linkage, and the pyranoside form of fructose were analyzed as follows. There was one methylene carbon not assigned yet, which was estimated as the C-1 of Fru (Fig. 3 (a)). Its protons showed HMBC correlations to methine carbon at δ_C 69.29 and the only one quaternary carbon (δ_C 101.52) (Fig. 3 (b)), which was assigned as C-3 and C-2 of Fru, respectively. The C-2 of Fru was also correlated to H-6 of Glc and H-6 of Fru (Fig. 3 (b)). These results revealed that saccharide 2 had a fructopyranoside residue and a Fru₂ 2 \rightarrow 6 Glc linkage, and all 1H and ^{13}C NMR signals were assigned. Due to strong coupling between H-3 and H-4 in β -Glc, H-3 and H-4 in Fru, H-5 and H-6 in Fru, and H-5 and H-6 in α -Glc, these couplings could not be analyzed in the first order. The δ_C values of Fru indicated its β anomer form, by comparing those of α and β




Fig. 3 Part of HSQC (A) and HMBC (B) spectra of saccharide 2.




Fig. 4 Structures of novel saccharides containing fructopyranoside isolated from fermented beverage of plant extract.

form of methyl-D-fructopyranoside (Sinclair 1988).

From these findings, the fructose residue of the non-reducing terminal of this saccharide was of a pyranose form, and saccharide 2 discovered in the fermented beverage of plant extracts was confirmed to be a novel saccharide, *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-D-glucopyranose (Fig. 4, Appendix 1).

Saccharides 10 and 14 were similarly identified as follows by examination of partial acid hydrolysates, GLC analysis of methanolysates and NMR measurement. Partial acid hydrolysates of saccharides 10 and 14 were liberated to glucose, fructose and laminaribiose. Saccharides 10 and 14 also had a fructose residue of the pyranose form. The methanolysate of permethylated saccharide 10 exhibited six peaks (Table 1), four of which corresponded to the α/β isomers of methyl 2,3,4-tri-*O*-methyl-D-glucopyranoside (t_R , 2.51 and 3.63) and methyl 2,4,6-tri-*O*-methyl-D-glucopyranoside (t_R , 3.26 and 4.77). The other two peaks corresponded to the α/β isomers methyl 1,3,4,5-tetra-*O*-methyl-D-fructopyranoside (t_R , 0.97 and 1.43).

The methanolysate of permethylated saccharide 14 exhibited three peaks, which corresponded to methyl 2,3,4,6-tetra-*O*-methyl-D-glucopyranoside (t_R , 1.04 and 1.46) and methyl 1,3,4,5-tetra-*O*-methyl-D-fructopyranoside (t_R , 0.92 and 1.46). No peaks corresponding to methyl 1,3,4,6-tetra-*O*-methyl-D-fructofuranoside were detected. The methanolysate of permethylated saccharide 14 gave two other peaks that were confirmed to be methyl 2,4-di-*O*-methyl-D-glucopyranoside from the retention time and pattern of fragmentation with GC-MS.

The 1H and ^{13}C NMR signals of each saccharide were assigned using 2D-NMR including COSY, HSQC, HSQC-TOCSY, CH_2 -HSQC-TOCSY, and CT-HMBC experiments (Appendix 4 and 7). Saccharides 10 and 14 were determined as new saccharides, *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-*O*- β -D-glucopyranosyl-(1 \rightarrow 3)-D-glucopyranose and *O*- β -D-fructopyranosyl-(2 \rightarrow 6)-[*O*- β -D-glucopyranosyl-(1 \rightarrow 3)]-D-glucopyranose, respectively (Fig. 4). No saccharides containing fructopyranoside residues were found in natural resources excepting the saccharide in the fermented beverage of plant extract. Furthermore, synthesis of saccharides 2, 10, and 14 by fermentation of plant extracts was investigated by using the ABEE-method and these saccharides were produced during fermentation (Fig. 5).

Partial acid hydrolysate of saccharide 12 was liberated to glucose, fructose and sucrose, and that of saccharide 13 was liberated to galactose, glucose, fructose and sucrose.




Fig. 5 High performance liquid chromatogram of fermentation products. HPLC analysis of saccharides produced during fermentation was done by ABEE conversion method. (A) Plant extract was fermented for 0 days. (B) Plant extract was fermented for 180 days. G: glucose; F: fructose.




Fig. 6 Structures of novel saccharides isolated from fermented beverage of plant extract.

The methanolsate of permethylated saccharide 12 exhibited four peaks (Table 1) corresponding to methyl 2,3,4,6-tetra-*O*-methyl-D-glucopyranoside (t_R , 1.01 and 1.41) and methyl 3,4,6-tri-*O*-methyl-D-fructofuranoside (t_R , 2.64 and 3.94). The methanolsate of permethylated saccharide 13 also exhibited six peaks (Table 1) corresponding to methyl 2,3,4,6-tetra-*O*-methyl-D-glucopyranoside (t_R , 1.01 and 1.42), methyl 2,3,4,6-tetra-*O*-methyl-D-galactopyranoside (t_R , 1.74), and methyl 3,4,6-tri-*O*-methyl-D-fructofuranoside (t_R , 2.64 and 3.97).

From these findings as above, saccharides 12 and 13 were proved to be D-glucosyl-(1 \rightarrow 1)-D-fructosyl-(2 \leftrightarrow 1)-D-glucoside and D-galactosyl-(1 \rightarrow 1)-D-fructosyl-(2 \leftrightarrow 1)-D-glucoside, respectively. The ^1H and ^{13}C NMR signals of each saccharide were assigned using 2D-NMR including COSY, HSQC, HSQC-TOCSY, and HMBC experiments (Appendix 5 and 6).

The saccharide 12 and 13 were identified as new saccharides, *O*- β -D-glucopyranosyl-(1 \rightarrow 1)-*O*- β -D-fructofuranosyl-(2 \leftrightarrow 1)- α -D-glucopyranoside; 1^F - β -D-glucopyranosylsucrose and *O*- β -D-galactopyranosyl-(1 \rightarrow 1)-*O*- β -D-fructofuranosyl-(2 \leftrightarrow 1)- α -D-glucopyranoside; 1^F - β -D-galactopyranosylsucrose, respectively (Fig. 6). Although, melezitose, 1^F - α -galactosylsucrose and 6^F - β -galactosylsucrose, had already been reported (Pazur *et al.* 1958; Courtois and




Fig. 7 High performance liquid chromatogram of fermentation products. Analysis of saccharides produced during fermentation was done by HPAEC. (A) Plant extract was fermented for 0 days. (B) Plant extract was fermented for 180 days. The beverage (100 mL) fermented for 0 or 180 days was added with charcoal (10 g), stirred for 3 h and filtered. The charcoal was extracted with 30% ethanol (500 mL) three times. The ethanol extracts were combined, concentrated to dryness and solubilized with one mL of distilled water. The sugar solution was analyzed by HPAEC.




Fig. 8 Structures of novel trisaccharides isolated from fermented beverage of plant extract.

Ariyoshi 1960), 1^F - β -D-glucopyranosylsucrose and 1^F - β -D-galactopyranosylsucrose were found in natural resources for the first time. Synthesis of the saccharides by fermentation of plant extract was investigated using HPAEC.

On the other hand, acid hydrolysates of saccharides 8 and 9 were liberated to glucose and fructose, and that of saccharide 16 was liberated to glucose. The methanolsate of permethylated saccharide 8 exhibited six peaks (Table 1) corresponding to methyl 2,3,4,6-tetra-*O*-methyl-D-glucopyranoside (t_R , 0.94 and 1.48), methyl 2,4,6-tri-*O*-methyl-D-glucopyranoside (t_R , 3.27 and 4.81) and methyl 1,3,4,6-tetra-*O*-methyl-D-fructofuranoside (t_R , 1.06 and 1.32). The methanolsate of permethylated saccharide 9 also exhibited six peaks (Table 1) corresponding to methyl 2,3,4-tri-*O*-methyl-D-glucopyranoside (t_R , 2.58 and 3.59), methyl 2,4,6-tri-*O*-methyl-D-glucopyranoside (t_R , 3.22 and 4.73), and methyl 1,3,4,6-tetra-*O*-methyl-D-fructofuranoside (t_R , 1.07

and 1.29). On the other hand, the methanolysate of permethylated saccharide **16** exhibited two peaks (Table 1) corresponding to methyl 2,3,4,6-tetra-*O*-methyl-*D*-glucopyranoside (t_R , 0.97 and 1.47). GC-MS analysis of the retention times and fragmentation patterns of methyl glucosides showed two peaks (10.08 and 10.21 min) from the methanolysate of permethylated saccharide **16** to be methyl 3,6-di-*O*-methyl-*D*-glucopyranoside. The ^1H and ^{13}C NMR signals of each saccharide were assigned using 2D-NMR including COSY, HSQC, HSQC-TOCSY, CH_2 -selected E-HSQC-TOCSY, HMBC and CT-HMBC experiments (Appendix 2, 3 and 8).

The saccharides **8**, **9** and **16** were *O*- β -*D*-fructofuranosyl-(2 \rightarrow 1)-[*O*- β -*D*-glucopyranosyl-(1 \rightarrow 3)]-*D*-glucopyranoside, *O*- β -*D*-fructofuranosyl-(2 \rightarrow 6)-*O*- β -*D*-glucopyranosyl-(1 \rightarrow 3)-*D*-glucopyranose and *O*- β -*D*-glucopyranosyl-(1 \rightarrow 2)-[*O*- β -*D*-glucopyranosyl-(1 \rightarrow 4)]-*D*-glucopyranose, respectively (Fig. 8), and were also novel oligosaccharides.

Synthesis of saccharides by fermentation of plant extracts was investigated using HPAEC. Almost all monosaccharides were removed from the fermented and unfermented beverages of plant extracts by the batch method with charcoal. The saccharides **8**, **9**, **12**, **13** and **16** were observed in the fermented beverage, but not in the unfermented one. Therefore, these five saccharides were confirmed to be produced during fermentation of plant extracts (Fig. 7).

Characteristics of saccharide 2, *O*- β -*D*-fructopyranosyl-(2 \rightarrow 6)-*D*-glucopyranose (Fp2-6G)

The characteristics of Fp2-6G were investigated. Sweetness was 0.2-fold compared with that of sucrose. Cariogenicity was performed according to the method of Miyamura (1973). As shown in Fig. 9, Fp2-6G was a non-cariogenic sugar because *S. mutans* and oral bacteria produce no acid. Furthermore, Fp2-6G was hydrolyzed by pig pancreatin and rat intestinal enzymes 3.5 and 12.2% compared with starch and sucrose, respectively, and the results showed that Fp2-6G had low digestibility. LB broth was used as basal medium for testing utilization of the saccharides by intestinal bacteria. *D*-Glucose, sucrose, lactose, melibiose, palatinose, turanose, raffinose, 1-kestose, or Fp2-6G was added to the LB medium at a final concentration of 0.5% (w/v). After incubation at 37°C for 72 h under anaerobic condition, bacterial growth was measured by pH analysis of the medium. The symbols “+++”, “++”, “+”, “±” and “-” show the medium pH of 4.0-4.5, 4.5-5.0, 5.0-5.5, 5.5-6.0 and 6.0-7.0, respectively. *Bifidobacteria* and *Lactobacilli* are beneficial to both nutrition and health of human and animals, while some intestinal bacteria such as *Escherichia coli*, *Enterobacter cloacae* and *Enterococcus faecalis* are detrimental intestinal bacteria.

Under the normal condition of *Bifidobacteria* growth, the pH of the medium added no saccharide (control), Fp2-6G, 1-kestose, raffinose, turanose, palatinose, melibiose, lactose, sucrose or glucose was 6.51-6.68, 5.47-6.14, 4.08-4.40, 4.03-4.18, 4.13-6.05, 4.22-5.12, 4.00-4.16, 4.02-4.50,




Fig. 9 Cariogenicity of Fp2-6G. A 0.5 mL aliquot of 1% (w/v) Fp2-6G, palatinose, glucose or distilled water was added to a mixture of 1.5 mL of fresh saliva or culture of *S. mutans*, and 0.5 mL of a brain heart infusion broth. (A) *Streptococcus mutans* JCM 5705. (B) Human saliva. Symbols: ○, Distilled water; □, Sucrose; ◇, Palatinose; ●, Fp2-6G.

4.12-4.94 or 4.00-4.40, respectively. On the other hand, Fp2-6G as well as 1-kestose and turanose were not fermented by *Enterobacter cloacae*, *Escherichia coli* and *Enterococcus faecalis*. Moreover, Fp2-6G was selectively used by four beneficial bacteria strains of *Bifidobacteria* (Table 2).

SUPPLEMENTATION

NMR-chemical shifts of saccharides (saccharides **2**, **8**, **9**, **10**, **12**, **13**, **14**, and **16**) are summarized as appendixes.

ACKNOWLEDGEMENTS

We thank Prof. J. Kawabata and Dr. E. Fukushi (Hokkaido University, Sapporo) for measuring the NMR spectra.

REFERENCES

- Aue WP, Batrholdi E, Ernst RR (1975) Two-dimensional spectroscopy. Application to nuclear magnetic resonance. *The Journal of Chemical Physics* **64**, 2229-2246
- Bax A, Summers MF (1986) ^1H and ^{13}C assignments from sensitivity-enhanced detection of heteronuclear multiple quantum NMR. *Journal of the American Chemical Society* **108**, 2093-2094
- Ciucanu I, Costello CE (2003) Elimination of oxidative degradation during the per-*O*-methylation of carbohydrates. *Journal of the American Chemical Society* **125**, 16213-16219
- Courtois JE, Ariyoshi U (1960) Saccharose galactosides from the roots of *Cucubalus baccifer* (Caryophyllaceae). Study of their structure. *Bulletin de la Société de Chimie Biologique* **42**, 737-751
- Domke T (1991) A new method to distinguish between direct and remote signals in proton-relayed X, H correlations. *Journal of Magnetic Resonance* **95**, 174-177
- Fujimoto H, Nishida H, Ajisaka K (1988) Enzymatic syntheses of glucobioses by a condensation reaction with α -glucosidase, β -glucosidases and glucoamylase. *Agricultural and Biological Chemistry* **52**, 1345-1351
- Hakomori S (1964) A rapid permethylation of glycolipid and polysaccharide catalyzed by methylsulfonyl carbanion in dimethylsulfoxide. *Journal of Biochemistry* **55**, 208
- Hurd RE, John BK (1991) Gradient-enhanced proton-detected heteronuclear multiple-quantum coherence spectroscopy. *Journal of Magnetic Resonance* **91**, 648-653

Table 2 Utilization of Fp2-6G and several other saccharides by some human intestinal bacteria.

Bacterial species	Fp2-6G	1-Kestose	Raffinose	Turanose	Palatinose	Melibiose	Lactose	Sucrose	Glucose
<i>Bifidobacterium adolescentis</i> 2793	+	+++	+++	-	+++	+++	+++	+++	+++
<i>B. bifidum</i> 2777	±	+++	+++	-	++	+++	+++	+++	+++
<i>B. breve</i> 2776	-	+++	+++	++	+	+++	+++	+++	+++
<i>B. infantis</i> 2775	±	+++	+++	++	++	+++	+++	+++	+++
<i>B. longum</i> 2778	+	+++	+++	+++	+++	+++	+++	++	+++
<i>Lactobacillus acidophilus</i> 2243	-	+++	+	-	+++	+++	+++	+++	+++
<i>L. casei</i> 2036	-	+	++	-	±	+++	+++	++	+++
<i>L. fermentum</i> 2046	-	±	++	-	-	+++	+++	++	+++
<i>Enterobacter cloacae</i> 1180	-	-	++	-	±	+++	++	++	+++
<i>Escherichia coli</i> 1099	-	-	-	-	-	±	+	++	++
<i>Enterococcus faecalis</i> 2048	-	-	-	-	±	++	++	++	+++
<i>Clostridium perfringens</i> 1211	-	+	±	-	-	++	++	++	++

The symbols “+++”, “++”, “+”, “±” and “-” show pH 4.0-4.5, 4.5-5.0, 5.0-5.5, 5.5-6.0 and 6.0-7.0, respectively.

- Johnson DC (1986) Carbohydrate detection gains potential. *Nature* 321, 451-452
- Kawamura S, Watanabe T, Abe N, Matsuda K (1971) Transglucosylation products from β -linked glucobiose with crude glucoamylase from *endomyces* sp. *The Society for Bioscience and Bioengineering* 49, 725-732
- Kawazoe N, Okada H, Fukushi E, Yamamori A, Onodera S, Kawabata J, Shiomi N (2008a) Two novel oligosaccharides isolated from a beverage produced by fermentation of a plant extract. *Carbohydrate Research* 343, 549-554
- Kawazoe N, Okada H, Fukushi E, Yamamori A, Onodera S, Kawabata J, Shiomi N (2008b) Structural analysis of two trisaccharides isolated from fermented beverage of plant extract. *Open Glycoscience* 1, 25-30
- Kobayashi I, Yasuno S, Hashimoto H, Ikura K, Nakano H, Kitahata S (2002) Analysis of positional isomers of α -linked glucooligosaccharides by *p*-aminobenzoic ethyl ester-conversion method. *Journal of Applied Glycoscience* 49, 461-468
- Kohmoto T, Fukui F, Takaku H, Machida Y, Arai M, Mitsuoka T (1988) Effect of isomalto-oligosaccharides on human fecal flora. *Bifidobacteria Microflora* 7, 61-69
- Miyamura E (1973) pH Analysis of the bacterial fermentation of sugar alcohol, maltitol and lactitol, in human whole saliva. *Shigaku* 60, 717-731
- Murosaki S, Muroyama K, Yamamoto Y, Kusaka H, Liu T, Yoshikai Y (1999) Immunopotentiating activity of nigerooligosaccharides for the T helper 1-like immune response in mice. *Bioscience, Biotechnology and Biochemistry* 63, 373-378
- Murosaki S, Muroyama K, Yamamoto Y, Liu T, Yoshikai Y (2002) Nigerooligosaccharides augments natural killer activity of hepatic mononuclear cells in mice. *International Immunopharmacology* 2, 151-159
- Okada H, Fukushi E, Onodera S, Nishimoto T, Kawabata J, Kikuchi M, Shiomi N (2003) Synthesis and structural analysis of five novel oligosaccharides prepared by glucosyltransfer from β -D-glucose 1-phosphate to isokestose and nystose using *Thermoanaerobacter brockii* kojibiose phosphorylase. *Carbohydrate Research* 338, 879-885
- Okada H, Fukushi E, Yamamori A, Kawazoe N, Onodera S, Kawabata J, Shiomi N (2006) Structural analysis of a novel saccharide isolated from fermented beverage of plant extract. *Carbohydrate Research* 341, 925-929
- Okada H, Kudoh K, Fukushi E, Onodera S, Kawabata J, Shiomi N (2005) Antioxidative activity and protective effect of fermented plant extract on ethanol-induced damage to rat gastric mucosa. *Journal of the Japanese Society of Nutrition and Food Science* 58, 209-215
- Okada H, Fukushi E, Yamamori A, Kawazoe N, Onodera S, Kawabata J, Shiomi N (2009) Structural analysis of three novel trisaccharides isolated from the fermented beverage of plant extracts. *Chemistry Central Journal* 3, 8
- Pachler KGR, Wessels PL (1973) Selective Population Inversion (SPI). A pulsed double resonance method in FT NMR spectroscopy equivalent to INDROR. *Journal of Magnetic Resonance* 12, 337-339
- Pazur JH, Marsh JM, Tipton CL (1958) Cosubstrate specificity of the transgalactosylase of *Saccharomyces fragilis*. *Journal of Biological Chemistry* 233, 277-279
- Rastall RA, Pikett SF, Adlard MW, Bucke C (1992) Synthesis of oligosaccharides by reversal of a fungal β -glucanase. *Biotechnology Letters* 14, 373-378
- Rocklin RD, Pohl CA (1983) Determination of carbohydrate by anion exchange chromatography with pulse amperometric detection. *Journal of Liquid Chromatography* 6, 1877-1590
- Shiomi N, Onodera S, Chatterton NJ, Harrison PA (1991) Separation of fructooligosaccharide isomers by anion-exchange chromatography. *Agricultural and Biological Chemistry* 55, 1427-1428
- Sinclair HB (1988) Two methyl anhydro-D-fructopyranosides prepared from D-mannitol. *Carbohydrate Research* 181, 115-123
- Takeda H, Sato K, Kinoshita S, Sasaki H (1994) Production of 1-kestose by *Scopulariopsis brevicaulis* sp. *Journal of Fermentation and Bioengineering* 77, 386-389
- Uzawa J, Yoshida S (2004) A new selective population transfer experiment using a double pulsed field gradient spin-echo. *Magnetic Resonance in Chemistry* 42, 1046-1048
- Villa TG, Phaff HJ (1979) A method for preparing laminaribiose by using yeast endo-(1 \rightarrow 3)- β -D-glucanase. *Carbohydrate Research* 74, 369-370
- von Kienlin M, Moonen CTW, von der Toorn A, van Zijl PCM (1991) Rapid recording of solvent-suppressed 2D COSY spectra with inherent quadrupole detection using pulsed field gradients. *Journal of Magnetic Resonance* 93, 423-429
- Willker W, Leibfritz D, Kerssebaum R, Bermel W (1993) Gradient selection in inverse heteronuclear correlation spectroscopy. *Magnetic Resonance in Chemistry* 31, 287-292
- Yamamori A, Fukushi E, Onodera S, Kawabata J, Shiomi N (2002) NMR analysis of mono- and difructosyllactosucrose synthesized by 1F -fructosyltransferase purified from roots of asparagus (*Asparagus officinalis* L.). *Magnetic Resonance in Chemistry* 40, 541-544
- Yasuno S, Murata T, Kokubo K, Yamaguchi T, Kamei M (1997) Two-mode analysis by high-performance liquid chromatography of *p*-aminobenzoic ethyl ester-derivatized monosaccharides. *Bioscience, Biotechnology and Biochemistry* 61, 1944-1946

Appendix 1 1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 2.

		δ_C	δ_H		J_{HH}
Fru _p	1	62.19	3.76	d	11.7
			3.79	d	11.7
	2	101.52			
	3	69.29	3.91	m ^b	
	4	70.34	3.91	m	
	5	70.00	3.98	br s ^c	
β Glc	6	64.83	3.91	m	
			3.70	m	
	1	96.72	4.61	d	7.7
	2	74.94	3.22	dd	9.4, 7.7
	3	76.31	3.45	m	
	4	70.34	3.45	m	
α Glc	5	75.88	3.56	dd	9.3, 5.7
			3.65	dd	10.8, 5.7
	6	61.20	3.88	d	10.8
	1	92.87	5.21	d	3.7
	2	72.24	3.52	dd	9.3, 3.7
	3	73.39	3.68	dd	9.8, 9.3
	4	70.51	3.39	dd	9.8, 9.3
			3.92	m	
	5	71.62	3.92	m	
	6	61.24	3.67	m	
			3.85	m	

^a Chemical shifts of 1H (δ_H) and ^{13}C (δ_C) in ppm were determined relatively to the external standard of sodium [2,2,3,3- 2H_4]-3-(trimethylsilyl) propanoate in D₂O (δ_H 0.00 ppm) and 1,4-dioxane (δ_C 67.40 ppm) in D₂O, respectively.

^b m: multiplet.

^c br s: broad singlet

Appendix 2 1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 8.

		δ_C	δ_H		J_{HH}
Fru _f	1	62.02	3.79	d	12.6
			3.69	d	12.6
	2	105.47			
	3	76.61	4.29	d	8.6
	4	74.57	4.14	dd	8.6, 8.6
	5	82.47	3.94	ddd	8.6, 6.4, 3.2
Glc	6	63.04	3.82	m ^b	
			3.74	dd	12.6, 4.4
	1	95.20	4.90	d	7.9
	2	73.44	3.55	dd	9.8, 7.9
	3	85.15	3.80	dd	9.8, 7.1
	4	68.87	3.52	m	
Glc'	5	76.11	3.51	m	
			3.89	m	
	6	61.33	3.74	dd	12.6, 4.4
	1	103.57	4.77	d	7.4
	2	74.29	3.36	dd	9.8, 7.4
	3	76.37	3.54	dd	9.8, 8.9
	4	70.42	3.42	dd	9.5, 8.9
			3.50	ddd	9.5, 5.7, 1.1
	5	76.81	3.93	m	
6	61.53	3.74	dd	12.3, 5.7	

^a Chemical shifts of 1H (δ_H) and ^{13}C (δ_C) in ppm were determined relatively to the external standard of sodium [2,2,3,3- 2H_4]-3-(trimethylsilyl) propanoate in D₂O (δ_H 0.00 ppm) and 1,4-dioxane (δ_C 67.40 ppm) in D₂O, respectively.

^b m: multiplet.

Appendix 3 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 9.

		δ_{C}	δ_{H}		J_{HH}			δ_{C}	δ_{H}		J_{HH}	
Fru f'	1	60.95	3.76	d	12.3	Fru f'	1					
			3.68	d	12.3							
	2	104.53					2					
	3	^g 77.75	^c 4.17	d	8.1		3	^g 77.79	^c 4.17	d	8.2	
	4	75.28	^d 4.11	dd	8.1, 7.8		4		^d 4.12	dd	8.2, 7.9	
	5	81.93	3.87	m ^b			5					
	6	^h 63.11	3.83	m		6	^h 63.14					
			3.67	dd	12.3, 6.4							
Glc	1	104.04	4.68	d	7.9	Glc	1	104.12	4.67	d	7.9	
	2	74.08	3.39	dd	9.5, 7.9		2	74.10	3.40	dd	9.6, 7.9	
	3	76.24	^e 3.54	dd	9.5, 8.5		3		^e 3.54	dd	9.6, 8.9	
	4	ⁱ 70.59	3.46	dd	10.1, 8.5		4	ⁱ 70.64				
	5	75.83	3.62	m			5					
	6	61.72	^f 4.08	dd	10.8, 4.4		6		^f 4.07	dd	10.8, 4.4	
			3.70	dd	10.8, 6.4							
β Glc'	1	96.45	4.70	d	8.1	α Glc'	1	92.74	5.26	d	3.9	
	2	74.15	3.44	dd	9.8, 8.1		2	71.33	3.73	dd	9.6, 3.9	
	3	87.14	3.72	dd	9.8, 7.1		3	84.95	3.89	dd	9.6, 9.2	
	4	^j 69.22	3.53	m			4	^j 69.19	3.53	m		
	5	76.27	3.53	m			5	71.91	3.89	m		
	6	61.54	3.90	m			6	61.36	3.84	dd	12.3, 2.2	
			3.73	dd	12.1, 4.7			3.77	m			

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.

^{c-j} interchangeable.

Appendix 4 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 10.

		δ_{C}	δ_{H}		J_{HH}
Fru p	1	62.18	3.79	m ^b	
			3.79	m	
	2	101.58			
	3	69.50	3.89	m	
	4	70.36	3.90	m	
	5	69.95	3.99	dd	3.2, 1.5
	6	64.86	3.87	m	
			3.72	m	
Glc'	1	103.95	4.67	d	7.7
	2	74.08	3.38	dd	9.8, 7.7
	3	76.19	3.53	m	
	4	70.45	3.47	m	
	5	75.80	3.63	m	
	6	61.20	3.93	m	
			3.67	m	
β Glc	1	96.44	4.68	d	8.0
	2	74.11	3.42	dd	9.4, 8.0
	3	87.02	3.69	m	
	4	69.16	3.50	m	
	5	76.26	3.50	m	
	6	61.51	3.88	m	
			3.72	m	
α Glc	1	92.71	5.25	d	3.7
	2	71.27	3.72	dd	9.6, 3.7
	3	84.83	3.85		
	4	69.13	3.50		
	5	71.92	3.87		
	6	61.34	3.82		
			3.76		

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.

Appendix 5 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 12.

		δ_{C}	δ_{H}		J_{HH}
α Glc	1	93.32	5.43	d	3.9
	2	71.96	3.54	dd	10.5, 3.9
	3	73.44	3.74	dd	10.5, 9.2
	4	70.09	3.46	dd	10.0, 9.2
	5	73.27	3.84	ddd	10.0, 3.7, 2.7
	6	60.98	3.81	m ^b	
Fru f'	1	69.76	4.05	d	11.1
			3.80	d	11.1
	2	103.82			
	3	77.29	4.31	d	8.5
	4	74.53	4.06	dd	8.9, 8.5
	5	82.14	3.91	ddd	8.9, 6.4, 4.2
	6	63.02	3.81	m	
β Glc	1	103.60	4.51	d	8.0
	2	73.88	3.32	dd	9.2, 8.0
	3	76.44	3.50	dd	9.2, 9.2
	4	70.42	3.40	dd	9.8, 9.2
	5	76.82	3.45	ddd	9.8, 5.7, 2.0
	6	61.52	3.91	dd	12.4, 2.0
			3.73	dd	12.4, 5.7

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.

Appendix 6 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 13.

		δ_{C}	δ_{H}		J_{HH}
αGlc	1	93.36	5.44	d	3.9
	2	71.99	3.55	dd	10.1, 3.9
	3	73.45	3.75	dd	10.1, 9.3
	4	70.11	3.47	dd	9.9, 9.3
	5	73.29	3.85	ddd	9.9, 3.5, 3.0
	6	60.99	3.81	m^b	
Fru f	1	69.81	4.07	d	11.2
			3.80	d	11.2
	2	104.19			
	3	77.43	4.33	d	8.9
	4	74.56	4.06	dd	8.9, 8.4
	5	82.13	3.92	ddd	8.4, 6.4, 2.7
βGal	6	63.03	3.81	m	
	1	104.19	4.44	d	7.9
	2	71.57	3.56	dd	9.6, 7.9
	3	73.48	3.66	dd	9.6, 3.4
	4	69.50	3.93	d	3.4
	5	76.10	3.70	dd	8.4, 4.6
	6	61.85	3.80	dd	12.1, 8.4
			3.78	dd	12.1, 4.6

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.

Appendix 7 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 14.

		δ_{C}	δ_{H}		J_{HH}
Fru p	1	62.16	3.79		
			3.79	m^b	
	2	101.55			
	3	69.28	3.92	m	
	4	70.33	3.92	m	
	5	69.95	3.99	br s ^c	
	6	64.86	3.93	m	
Glc'			3.72	m	
	1	103.58	4.74	d	8.0
	2	74.26	3.35	dd	9.4, 8.0
	3	76.37	3.52	dd	9.4, 8.7
	4	70.39	3.41	dd	8.9, 9.7
	5	76.83	3.47	m	
βGlc	6	61.52	3.92	m	
			3.72	m	
	1	96.50	4.66	d	8.1
	2	74.63	3.43	dd	8.3, 8.1
	3	85.03	3.73	dd	8.3, 8.3
	4	68.91	3.56	m	
αGlc	5	75.57	3.59	m	
	6	61.23	3.90	m	
			3.69	m	
	1	92.79	5.23	d	3.6
	2	71.81	3.73	dd	9.6, 3.6
	3	82.83	3.90	dd	9.6, 9.6
	4	69.03	3.55		
	5	71.46	3.96		
	6	61.27	3.87		
			3.69		

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.

^c br s: broad singlet.

Appendix 8 ^1H and ^{13}C NMR spectral data (δ^a in ppm, J in Hz) of saccharide 16.

		δ_{C}	δ_{H}		J_{HH}
αGlc	1	92.35	5.43	d	4.0
	2	81.12	3.69	dd	9.7, 4.0
	3	71.22	3.96	dd	9.7, 9.7
	4	79.09	3.69	dd	10.1, 9.7
	5	70.70	3.95	ddd	10.1, 6.9, 1.0
	6	60.67	3.86	m^b	
Glc'	1	104.74	4.63	d	7.6
	2	74.11	3.35	dd	9.6, 7.6
	3	76.40	3.49	m	
	4	70.29	3.40	m	
	5	76.62	3.44	m	
	6	61.44	3.72	dd	12.3, 5.6
Glc''			3.89	dd	12.3, 1.2
	1	103.26	4.51	d	7.6
	2	73.98	3.31	dd	9.6, 7.6
	3	76.33	3.50	m	
	4	70.29	3.41	m	
	5	76.86	3.48	m	
βGlc	6	61.44	3.73	dd	12.3, 6.2
			3.91	dd	12.3, 1.2
	1	95.32	4.73	d	8.0
	2	81.72	3.55	dd	9.1, 8.0
	3	75.39	3.83	dd	9.4, 9.1
	4	79.09	3.68	dd	9.6, 9.4
Glc''	5	75.59	3.59	m	
	6	60.87	3.81	dd	11.7, 5.2
			3.95	dd	11.7, 1.2
	1	103.51	4.80	d	7.6
	2		3.31	dd	9.6, 7.6
	3				
Glc''	4				
	5				
	6		3.70	dd	12.2, 6.5
			3.92	dd	12.2, 1.2
	1	103.32	4.50	d	7.8
	2		3.30	dd	9.6, 7.8
			3.72	dd	12.3, 6.2

^a Chemical shifts of ^1H (δ_{H}) and ^{13}C (δ_{C}) in ppm were determined relatively to the external standard of sodium [2,2,3,3- $^2\text{H}_4$]-3-(trimethylsilyl) propanoate in D_2O (δ_{H} 0.00 ppm) and 1,4-dioxane (δ_{C} 67.40 ppm) in D_2O , respectively.

^b m: multiplet.