

目 次

村山三郎・小阪進一・山下憲之：アルファルファの品種生態に関する研究 3. 播種密度がアルファルファ品種の初期生育および 体内成分におよぼす影響	327
浅川満彦・神谷晴夫・大林正士：Studies on the Parasite Fauna of Insectivora II. Four new capillarid nematodes from the Japanese shrews, genera <i>Sorex</i> and <i>Crocidura</i>	335
浅川満彦：Genus <i>Heligmosomoides</i> HALL, 1916 (Heligmosomidae: Nematoda) from the Japanese wood mice, <i>Apodemus</i> spp. II. A review of the genus <i>Heligmosomoides</i> with the establishment of the phylogenetic lines of known species	349
浅川満彦・長谷川英男・福本真一郎：A Taxonomical Note on <i>Longistriata wolgaensis</i> sensu YAMAGUTI, 1954 (Nematoda: Heligmonellidae: Nippostrongylinae) from <i>Clethrionomys smithii</i>	367
荒木重美：On the Stability of a Skyrme-type Model in the D -dimensional Space	371
荒木重美：Note on the Berezin-Kostant Quantization	377
荒木重美：Structure of Vacua and Instability of Hopf Soliton in the Skyrme Model	389
荒木重美：Finite Dimensional Irreducible Representation of Noncompact Groups. II	395
松井幸夫・村山正道・西 誠治・飯沼昭博：Soybean blight caused by <i>Fusarium oxysporum</i> var. <i>redolens</i> and the production of Fusaric acid by the fungus	403

西埜 進・森田 茂・近藤誠司：人工乳主体時の石灰石給与が 離乳子牛の消化率と糞 pH に及ぼす影響	413
森田 茂・石村行弘・西埜 進：配合飼料と乾草の給与順序が 去勢牛の採食量におよぼす影響	425
野 英二・安宅一夫・原 宗之・楠崎 舜・井上錦次：乳牛への ルーメン・バイパスメチオニン製剤投与が乳量 および乳組成に及ぼす影響	435
守重哲郎・高橋清志・阿部 聡・黒沢 隆・其田三夫：Detection of Circulating Immune Complex in Dogs Infected with <i>Babesia gibsoni</i>	443
小山久一・鈴木 太：ゼラチンスライド法による家兎子宮および 腔内精子のアクロシン活性の観察	453

CONTENTS

MURAYAMA, S., KOSAKA, S. and YAMASHITA, N.: Studies on Varietal Ecology of Alfalfa Plants 3. Influence of Seeding Densities on Seedling Growth and Chemical Composition of Alfalfa Varieties	327
ASAKAWA, M., KAMIYA, H. and OHBAYASHI, M.: Studies on the Parasite Fauna of Insectivora II. Four new capillarid nematodes from the Japanese shrews, genera <i>Sorex</i> and <i>Crocidura</i>	335
ASAKAWA, M.: Genus <i>Heligmosomoides</i> HALL, 1916 (Heligmosomidae: Nematoda) from the Japanese wood mice, <i>Apodemus</i> spp. II. A review of the genus <i>Heligmosomoides</i> with the establishment of the phylogenetic lines of known species	349
ASAKAWA, M., HASEGAWA, H. and FUKUMOTO, S.: A Taxonomical Note on <i>Longistriata wolgaensis</i> sensu YAMAGUTI, 1954 (Nematoda: Heligmonellidae: Nippostrongylineae) from <i>Clethrionomys smithii</i>	367
ARAKI, S.: On the Stability of a Skyrme-type Model in the D -dimensional Space	371
ARAKI, S.: Note on the Berezin-Kostant Quantization	377
ARAKI, S.: Structure of Vacua and Instability of Hopf Soliton in the Skyrme Model	389
ARAKI, S.: Finite Dimensional Irreducible Representation of Noncompact Groups. II	395
MATSUI, Y., MURAYAMA, M., NISHI, S. and IHNUMA, A.: Soybean blight caused by <i>Fusarium oxysporum</i> var. <i>redolens</i> and the production of Fusaric acid by the fungus	403
NISHINO, S., MORITA, S. and KONDO, S.: Effect of Limestone Feeding to High Calf Starter Rations on Digestibility and Fecal pH in Weaned Calves	413

MORITA, S., ISHIMURA, Y. and NISHINO, S.: Effect of Feeding Order of Concentrate and Hay on Feed Intake in Steers	425
NO, E., ATAKU, K., HARA, M., NARASAKI, N. and INOUE, K.: Effect of Rumen-Bypass Methionine on Milk Production	435
MORISHIGE, T., TAKAHASHI, K., ABE, S., KUROSAWA, T. and SONODA, M.: Detection of Circulating Immune Complex in Dogs Infected with <i>Babesia gibsoni</i>	443
KOYAMA, H. and SUZUKI, H.: The Acrosin Activity in Rabbit Spermatozoa Recovered from Uterine Horn and Vagina by Gelatin Substrate Film Method	453