



# Use of *Aeromonas* spp. as General Indicators of Antimicrobial Susceptibility among Bacteria in Aquatic Environments in Thailand

Masaru Usui<sup>1</sup>, Chie Tagaki<sup>1</sup>, Akira Fukuda<sup>1</sup>, Torahiko Okubo<sup>1</sup>, Chanchai Boonla<sup>2</sup>, Satoru Suzuki<sup>3</sup>, Kanako Seki<sup>4</sup>, Hideshige Takada<sup>4</sup> and Yutaka Tamura<sup>1\*</sup>

<sup>1</sup> School of Veterinary Medicine, Rakuno Gakuen University, Ebetsu, Japan, <sup>2</sup> Departments of Biochemistry, Chulalongkorn University, Bangkok, Thailand, <sup>3</sup> Center for Marine Environmental Studies, Ehime University, Matsuyama, Japan, <sup>4</sup> Laboratory of Organic Geochemistry, Tokyo University of Agriculture and Technology, Fuchu, Japan

## OPEN ACCESS

### Edited by:

David W. Graham,  
Newcastle University, UK

### Reviewed by:

Tanel X. Tenson,  
University of Tartu, Estonia  
Maria Ines Zanoli Sato,  
CETESB – Companhia Ambiental do  
Estado de Sao Paulo, Brazil

### \*Correspondence:

Yutaka Tamura  
tamuray@rakuno.ac.jp

### Specialty section:

This article was submitted to  
Antimicrobials, Resistance  
and Chemotherapy,  
a section of the journal  
Frontiers in Microbiology

**Received:** 18 March 2016

**Accepted:** 28 April 2016

**Published:** 12 May 2016

### Citation:

Usui M, Tagaki C, Fukuda A, Okubo T,  
Boonla C, Suzuki S, Seki K, Takada H  
and Tamura Y (2016) Use  
of *Aeromonas* spp. as General  
Indicators of Antimicrobial  
Susceptibility among Bacteria  
in Aquatic Environments in Thailand.  
*Front. Microbiol.* 7:710.  
doi: 10.3389/fmicb.2016.00710

Antimicrobials are widely used, not only for treating human infections, but also for treatment of livestock and in fish farms. Human habitats in Southeastern Asian countries are located in close proximity to aquatic environments. As such, the human populations within these regions are at risk of exposure to antimicrobial resistant bacteria, and thereby disseminating antimicrobial resistance genes (ARGs). In this study, we collected water samples from 15 sites (5 sites in Chao Phraya River, 2 sites at the mouth of Chao Phraya River, 3 sites in Ta Chin River, and 5 sites at city canals) and 12 sites (6 sites at city canals; 2 sites at chicken farms; 2 sites at pig farms; and 2 samples from sites at pig farms, which were subsequently treated at a biogas plant) in Thailand in 2013 and 2014, respectively. In total, 117 *Aeromonas* spp. were isolated from the water samples, and these organisms exhibited various antimicrobial susceptibility profiles. Notably, there was a significant correlation between the environmental concentration of tetracyclines and the rates of tetracycline resistance in the isolated *Aeromonas* spp.; however, both the concentration and rates of tetracycline resistance in samples derived from pig farms were higher than those of samples harvested from other aquatic environments. These findings suggest that the high concentrations of antimicrobials observed in these aquatic environments likely select for ARGs. Furthermore, they indicate that *Aeromonas* spp. comprise an effective marker for monitoring antimicrobial resistance in aquatic environments.

**Keywords:** *Aeromonas* spp., antimicrobial resistance, aquatic environments, tetracycline resistance, indicator bacteria

## INTRODUCTION

The use of antimicrobials for treating/preventing bacterial infections is essential for promoting the health and welfare of humans and animals. As such, the emergence and spread of antimicrobial resistance is of global concern. To address these issues, the World Health Organization (WHO) and other international organizations require the development of programs for monitoring antimicrobial resistance in zoonotic bacteria, animal pathogens, and indicator bacteria derived from food-producing animals (Dehaumont, 2004; FAO/OIE/WHO, 2004).

Antimicrobials are widely used, not only for treating human infections, but also in livestock and fish farms. As a consequence, large quantities of these antimicrobials are released into aquatic environments via sewage or directly from livestock and fish farms, thereby providing a selective pressure for the development or acquisition of antimicrobial-resistant bacteria (ARB) and genes (ARGs) (Esteve et al., 2015). Indeed, aquatic environments have been shown to function as reservoirs of ARB and ARGs (Marti et al., 2014), and the transfer of plasmid-encoded ARGs from aquatic environments to humans comprises the greatest potential risk to human health (Smith et al., 1994). To date, however, there have been few reports regarding the monitoring of antimicrobial resistance in aquatic environments.

To efficiently monitor antimicrobial resistance, it is necessary to identify an effective general indicator bacterium within aquatic environments. For example, *Escherichia coli* is commonly used to monitor antimicrobial resistance in clinical settings and in the veterinary field (NARMS, 2012; National Veterinary Assay Laboratory Forestry and Fisheries, 2012); however, this organism is not always isolated from aquatic environments. In contrast, *Aeromonas* spp. are Gram-negative, waterborne organisms that are ubiquitous in most aquatic environments (Rhodes et al., 2000). We therefore selected *Aeromonas* spp. as a general model organism to inform on antimicrobial resistance in aquatic environments.

The spread and maintenance of ARB and ARGs in aquatic environments comprises a potential human health risk. In Southeastern Asian countries, including Thailand, human habitats are often located in close proximity to aquatic environments. Indeed, contamination of both humans and livestock animals with water-derived ARB has been demonstrated in tropical Asian countries (Shimizu et al., 2013). Likewise, a previous study detected contamination of individuals in Indochina with ARGs derived from aquatic environments (Suzuki and Hoa, 2012). Thailand is one of the most agriculturally productive countries in Southeast Asia, and livestock animals in this country are frequently administered antimicrobials to treat and prevent infections (Daniel et al., 2015). As a result, individuals in Thailand are likely at a greater risk of exposure to ARB and ARGs than those in other countries in this region. In this study, we first demonstrated the usefulness of *Aeromonas* spp. as general indicator bacteria for monitoring antimicrobial resistance. We subsequently utilized this organism to evaluate the current prevalence of resistance at multiple aquatic sites in Thailand. Finally, we examined putative correlations between the prevalence of ARB and ARGs and drug concentrations in rivers and livestock wastewater.

## MATERIALS AND METHODS

### Sample Collection

Grab water samples were collected in sterile 1-L bottles in Bangkok and in the province of Ratchaburi in September 2013 and 2014 (Figure 1). The characteristics of the sampling sites

are summarized in Supplementary Table S1. Specifically, in 2013, surface water samples were taken at 5 sites (RC 1, 2, 3, 4, and 5) on the Chao Phraya River and at 2 sites (RCM 1 and 2) at the mouth of Chao Phraya River. In addition, 3 sites (RT 1, 2, and 3) were sampled on the Ta Chin River and 5 samples were harvested at city canals (C 1, 2, 3, 4, and 5) in Bangkok. In 2014, 6 samples were taken at city canals (C 6, 7, 8, 9, 10, and 11) in Bangkok, and effluent was collected from chicken farms (FC 1 and 2) and pig farms (FP 1 and 2) in the province of Ratchaburi. The pig farm effluent was also treated at a biogas plant. Treated effluent is indicated as post-treatment FP 1 and 2 (FFP1 and FFP2). All samples were stored at 4°C after collection and analyzed within 12 h.

### Analysis of Antimicrobial Concentrations


To measure aquatic concentrations of antimicrobials, including sulfonamides and tetracyclines, water samples were analyzed using a liquid chromatograph (Accela, Thermo Scientific, Waltham, MA, USA) equipped with a tandem mass spectrometer (LC-MS/MS; Quantum Access, Thermo Scientific) after extraction using a solid-phase cartridge (Oasis HLB resin; Waters Corp., Milford, MA, USA). The analytical process was the same as that described by Segura et al. (2015).

### Bacterial Isolation

For isolation of *Aeromonas* spp. from water samples, deoxycholate-hydrogen sulfate-lactose (DHL) agar medium (Nissui Pharmaceutical, Tokyo, Japan) was inoculated with 100 µL of each respective sample and incubated at 37°C overnight. For each sample, a maximum of six colonies were identified as likely *Aeromonas* spp. based on colony morphology on DHL agar (pink colony) and were subjected to further analyses. Specifically, the isolates were screened for oxidase production and subjected to a variety of biochemical tests (API20E; BioMérieux, Marcy l'Etoile, France). Furthermore, the identity of each strain was verified by 16S rRNA gene sequencing (Marchesi et al., 1998).

### Antimicrobial Susceptibility Testing

The minimal inhibitory concentrations (MICs) of the following antimicrobials for each isolate were determined using the agar dilution method, according to the Clinical Laboratory Standards Institute (CLSI) guidelines (Clinical Laboratory Standards Institute, 2008): ampicillin, cefazolin, cefotaxime, kanamycin, tetracycline, sulfamethoxazole, nalidixic acid, and ciprofloxacin (all obtained from Sigma-Aldrich, St. Louis, MO, USA). The breakpoints for each antimicrobial were in accordance with CLSI guidelines (Clinical Laboratory Standards Institute, 2008), except for those of sulfamethoxazole, which are not defined by the CLSI guidelines. As such, we utilized the break points for sulfonamide for this antimicrobial. *E. coli* ATCC25922, *Staphylococcus aureus* ATCC29213, *Enterococcus faecalis* ATCC29212, and *Pseudomonas aeruginosa* ATCC27853 were used as quality control strains.



**FIGURE 1 | Map of sampling sites in Thailand.** RC 1–5 indicate Chao Phraya River sites; RCM 1 and 2 indicate sites at the mouth of Chao Phraya River; RT 1–3 indicate Ta Chin River sites; C 1–5 indicate the city canal sites sampled in 2013; C 6–11 indicate the city canal sites sampled in 2014; FC 1 and 2 and FP 1 and 2 indicate chicken and pig farm sites, respectively. Sources: [http://www.savgis.org/cv\\_EN.htm](http://www.savgis.org/cv_EN.htm).

## Characterization of ARGs

Each strain was evaluated for the presence of the *tetA*, *tetB*, *tetC*, *tetD*, *tetE*, and *tetG* genes via multiplex PCR, as previously described (Jin Jun et al., 2004). Meanwhile, PCR analysis was utilized to detect the *tetM* and *tetS* genes (Kim et al., 2004) and the *sul1*, *sul2*, and *sul3* genes (Hoa et al., 2008), respectively.

## Statistical Analysis

Correlations between the environmental concentration of each antimicrobial and both the rate of antimicrobial resistance and the prevalence of resistance genes among *Aeromonas* spp., respectively, were evaluated using Pearson's test. Differences were considered significant at  $p < 0.05$ .

## RESULTS

### Isolation of *Aeromonas* spp. from Water Samples

In total, 117 *Aeromonas* strains were isolated from 27 water samples. Specifically, 24, 9, 13, 13, 30, 9, 12, and 7 strains were isolated from riverine of Chao Phraya River, the mouth of Chao Phraya River, the riverine of Ta Chin River, the city canal, chicken farms, pig farms, and post-treatment pig farms, respectively (Table 1).

## Antimicrobial Resistance


The proportions of antimicrobial-resistant *Aeromonas* spp. are summarized in Figure 2. Nearly all isolates were resistant to ampicillin and cefazolin. Likewise, we frequently detected sulfamethoxazole- and nalidixic acid-resistant isolates. Notably, however, the rates of cefotaxime, kanamycin, tetracycline, and ciprofloxacin resistance were markedly higher among isolates derived from pig farms than among those obtained from other sites.

### Antimicrobial Resistance Genes (ARGs)

While the *tetA* gene was detected in multiple river and canal samples, the prevalence of this gene was highest among the pig farm isolates (75.0%; Table 1). Moreover, both the *tetM* and *sul3* genes (16.7 and 25%, respectively; Table 1) were detected only in isolates harvested from pig farms, while the *sul2* gene was detected in pig farm and canal (2014) isolates only (33.3 and 3.3%, respectively; Table 1). The *tetC* and *tetE* genes were the most frequently detected resistance genes among the river isolates (Table 1), and the *sul1* gene was detected in all water environments except for chicken farms (15.4 to 44.4%). Notably, the *tetC*, *tetG*, and *tetS* genes were not detected in any of the isolates examined in this study, while *tetB* was detected in only 1 of the 30 canal isolates from 2014 (3.3%).

**TABLE 1 | Prevalence of antimicrobial resistance genes (ARGs) among *Aeromonas* spp. isolated from aquatic environments in Thailand.**

Sampling sites	n	Tetracycline resistance genes					Sulfonamide resistance genes		
		tetA (%)	tetB (%)	tetC (%)	tetE (%)	tetM (%)	sul1 (%)	sul2 (%)	sul3 (%)
Chao Phraya River	24	4.2	0	8.3	29.1	0	37.5	0	0
Mouth of the Chao Phraya River	9	11.1	0	11.1	33.3	0	44.4	0	0
Ta Chin River	13	0	0	15.4	0	0	15.4	0	0
City Canal, 2013	13	7.7	0	7.7	7.7	0	30.8	0	0
City Canal, 2014	30	0	3.3	20	23.3	0	26.7	3.3	0
Chicken Farm	9	0	0	0	0	0	0	0	0
Pig Farm	12	75	0	0	0	16.7	25	33.3	25
Pig Farm, Post-treatment	7	28.5	0	0	0	14.3	28.6	42.9	14.3



## Relationship between Rates of Antimicrobial Resistance and Concentrations of Antimicrobials

The average concentrations of tetracyclines and sulfonamides at each sample site are summarized in **Table 2**. Subsequent statistical analyses detected significant correlations between the environmental concentration of tetracyclines and both the rate of tetracycline resistance and the prevalence of tetracycline resistance genes among *Aeromonas* spp. ( $R^2 = 0.63$  and  $R^2 = 0.57$ ,  $p = 0.019$  and  $p = 0.030$ , respectively; **Figure 3**). In contrast, there was no statistical correlation between the environmental concentration of sulfonamides and the rate of sulfamethoxazole resistance

or the prevalence of sulfonamide resistance genes among *Aeromonas* spp.

## DISCUSSION

In this study, *Aeromonas* spp. were frequently isolated from water samples. Notably, these isolates exhibited various resistance patterns against the antimicrobials tested, with the exception of ampicillin and cefazolin, to which nearly all strains were resistant. These findings are therefore consistent with those of a previous study demonstrating that *Aeromonas* spp. are endogenously resistant to these two antimicrobials (Cisar et al., 2014). Meanwhile, the cephalosporin-resistant *Aeromonas* spp.

**TABLE 2 | The average concentrations of tetracyclines and sulfonamides at the aquatic environments examined in this study.**

Sampling sites	n	Average concentrations (ng/L)	
		Tetracyclines	Sulfonamides
Chao Phraya River	5	<2	12.6 ± 5.0
Mouth of the Chao Phraya River	2	<2	13.7 ± 2.4
Ta Chin River	3	<2	33.7 ± 1.9
City Canal, 2013	5	<2	74.8 ± 54.8
City Canal, 2014	6	138.8 ± 123.2	111.5 ± 119.2
Chicken Farm	2	482.0 ± 503.5	ND
Pig Farm	2	43,500.0 ± 4,0527.1	48.0 ± 9.9
Pig Farm, Post-treatment	2	2,135.5 ± 365.6	12.5 ± 16.3

ND, not detected.

exhibited resistance against one of the latest generation of cephalosporins, cefotaxime. As such, monitoring of cefotaxime resistance in *Aeromonas* spp. would provide information regarding the resistance rates to latest generation cephalosporins among aquatic isolates. Furthermore, Yano et al. (2015) previously reported that *Aeromonas* spp. exhibited a greater variety of antimicrobial susceptibility profiles than other aquatic bacteria (Yano et al., 2015). Together, these results suggest that *Aeromonas* spp. comprise an effective general indicator organism for monitoring antimicrobial resistance in aquatic environments.


*Aeromonas* spp. were previously found to be as prevalent in humans as in environmental sources, indicating that these organisms could disseminate antimicrobial resistance from environmental to clinical environments (Sanchez-Cespedes et al., 2009). Meanwhile, yet-to-be cultured bacteria in aquatic environments can act as non-visible ARG reservoirs (Suzuki et al., 2015), and may be capable of transferring these ARGs to both *Aeromonas* spp. and other pathogenic aquatic bacteria. As such, monitoring the prevalence and dissemination of antimicrobial resistance in aquatic environments using *Aeromonas* spp. as an indicator organism is essential to minimize potential risks to human health.

In this study, we detected 40-fold higher concentrations of tetracyclines in pig farm wastewater than in river water samples. Indeed, in the two pig farms tested, chlortetracycline

was used as a feed additive for 15- to 20-week-old piglets (personal communications). In contrast, the concentrations of sulfonamides detected in this study were markedly lower than those previously detected on pig farms in Vietnam and South Africa (Le and Muneke, 2004; Hoa et al., 2011; Shimizu et al., 2013; Suzuki et al., 2015). These results suggest that while tetracyclines are currently used intensively, sulfonamides have begun to be phased out for agricultural use in Thailand.

We detected a significant correlation between the environmental concentration of tetracyclines and the rate of tetracycline resistance, but not between the concentration of sulfonamides and the rate of sulfonamide resistance among the *Aeromonas* strains isolated in this study. Our findings are therefore partially consistent with those of a previous study, which reported that high concentrations of antimicrobials (ciprofloxacin, ofloxacin, cefazolin, cefotaxime, sulfamethoxazole, and clarithromycin) were associated with high copy numbers of ARGs in hospital wastewater samples (Rodriguez-Mozaz et al., 2015). Indeed, these high concentrations of antimicrobials played an important role in the selection and maintenance of ARB and ARGs. These findings therefore suggest that aquatic environments, particularly those containing high concentrations of antimicrobials, act as reservoirs of ARB and ARGs. However, our sampling data were limited, especially in regard to sampling points associated with high concentrations of antimicrobials. As such, in future studies, a greater number of such sampling points is needed to obtain more reliable findings.

The *tetA* gene was the most frequently detected ARG among tetracycline-resistant *Aeromonas* isolates derived from pig farms. In a previous report, *tetA* was the most prevalent tetracycline-resistant gene among *E. coli* strains isolated from pig feces (Schwaiger et al., 2010), suggesting that the *tetA* gene of *Aeromonas* spp. isolated from pig farm wastewater originated from pig feces. Meanwhile, the *tetE* gene was the mostly frequently observed ARG among isolates derived from river and canal water. Notably, *tetE* has also been detected in aquatic sediments from aquaculture farms, despite the absence of tetracycline treatments (Tamminen et al., 2011). In most aquatic environments, antimicrobial concentrations are low [e.g., sub- to low- $\mu\text{g/L}$  levels, even in anthropogenically impacted waters



(Shimizu et al., 2013)]. However, long-term consistent exposure to sub-threshold doses of antimicrobials can result in increased rates of bacterial resistance (Harnisz, 2013). These results suggest that antimicrobial resistance genes (ARGs), including *tetE*, can be maintained in aquatic environments regardless of the concentration of the corresponding antimicrobial. Consistent with this conclusion, we recently demonstrated that multidrug resistance plasmids can be retained without selective pressure in viable but non-culturable cells in oligotrophic water environments (Bien et al., 2015).

In this study, treatment of pig farm samples at a biogas plant resulted in decreased rates of resistance to six antimicrobials, as well as reduced concentrations of antimicrobials. Likewise, a previous study demonstrated that biogas plant treatments decrease the prevalence of ARB and tetracyclines in cattle feces (Ihara et al., 2012). These results suggest that biogas plant treatments comprise an effective method for decreasing the concentration of certain antimicrobials and ARB in livestock wastewater.

## REFERENCES

- Bien, T. L., Sato-Takabe, Y., Ogo, M., Usui, M., and Suzuki, S. (2015). Persistence of multi-drug resistance plasmids in sterile water under very low concentrations of tetracycline. *Microbes Environ.* 30, 339–343. doi: 10.1264/jisme2.ME15122
- Cisar, C. R., Henderson, S. K., Askew, M. L., Risenhoover, H. G., McAndrews, C. R., Kennedy, S. D., et al. (2014). Antibiotic resistance in *Aeromonas* upstream and downstream of a water resource recovery facility. *Water Environ. Res.* 86, 835–843. doi: 10.2175/106143014X14062131177917
- CLSI Clinical and Laboratory Standards Institute (2008). *Performance Standards for Antimicrobial Disk Dilution Susceptibility Tests for Bacteria Isolated from Animals: Approved Standard-Third Edition Documents M31-A3*. Wayne, PA: CLSI.
- Daniel, D. S., Lee, S. M., Dykes, G. A., and Rahman, S. (2015). Public health risks of multiple-drug-resistant *Enterococcus* spp. in Southeast Asia. *Appl. Environ. Microbiol.* 81, 6090–6097. doi: 10.1128/AEM.01741-15
- Dehaumont, P. (2004). OIE International standards on antimicrobial resistance. *J. Vet. Med. B Infect. Dis. Vet. Public Health* 51, 411–414. doi: 10.1111/j.1439-0450.2004.00784.x
- Esteve, C., Alcaide, E., and Gimenez, M. J. (2015). Multidrug-resistant (MDR) *Aeromonas* recovered from the metropolitan area of Valencia (Spain): diseases spectrum and prevalence in the environment. *Eur. J. Clin. Microbiol. Infect. Dis.* 34, 137–145. doi: 10.1007/s10096-014-2210-z
- FAO/OIE/WHO (2004). *Joint FAO/OIE/WHO Expert Workshop on Non-Human Antimicrobial Usage and Antimicrobial Resistance: Scientific Assessment*. Available at <http://www.oie.int/doc/ged/D895.PDF>
- Harnisz, M. (2013). Total resistance of native bacteria as an indicator of changes in the water environment. *Environ. Pollut.* 174, 85–92. doi: 10.1016/j.envpol.2012.11.005
- Hoa, P. T., Managaki, S., Nakada, N., Takada, H., Shimizu, A., Anh, D. H., et al. (2011). Antibiotic contamination and occurrence of antibiotic-resistant bacteria in aquatic environments of northern Vietnam. *Sci. Total Environ.* 409, 2894–2901. doi: 10.1016/j.scitotenv.2011.04.030
- Hoa, P. T. P., Nonaka, L., Hung Viet, P., and Suzuki, S. (2008). Detection of the *sul1*, *sul2*, and *sul3* genes in sulfonamide-resistant bacteria from wastewater and shrimp ponds of north Vietnam. *Sci. Total Environ.* 405, 377–384. doi: 10.1016/j.scitotenv.2008.06.023
- Ihara, I., Yoshida, G., Toyoda, K., Iwasaki, M., and Umetsu, K. (2012). Environmental impacts of veterinary antibiotics and reduction of related risk by anaerobic digestion. *Soil Microorgan.* 60, 47–50.
- Jin Jun, L., Bum Jeong, J., Huh, M. D., Chung, J. K., Choi, D. L., Lee, C. H., et al. (2004). Detection of tetracycline-resistance determinants by multiplex polymerase chain reaction in *Edwardsiella tarda* isolated from fish farms in Korea. *Aquaculture* 240, 89–100. doi: 10.1016/j.aquaculture.2004.07.025
- Kim, S. R., Nonaka, L., and Suzuki, S. (2004). Occurrence of tetracycline resistance genes *tet(M)* and *tet(S)* in bacteria from marine aquaculture sites. *FEMS Microbiol. Lett.* 237, 147–156. doi: 10.1016/j.femsle.2004.06.026
- Le, T. X., and Munekage, Y. (2004). Residues of selected antibiotics in water and mud from shrimp ponds in mangrove areas in Viet Nam. *Mar. Pollut. Bull.* 49, 922–929. doi: 10.1016/j.marpolbul.2004.06.016
- Marchesi, J. R., Sato, T., Weightman, A. J., Martin, T. A., Fry, J. C., Hiom, S. J., et al. (1998). Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Appl. Environ. Microbiol.* 64, 795–799.
- Marti, E., Variatza, E., and Balcazar, J. L. (2014). The role of aquatic ecosystems as reservoirs of antibiotic resistance. *Trends Microbiol.* 22, 36–41. doi: 10.1016/j.tim.2013.11.001
- NARMS (2012). *NARMS Reports and Data*. Available at: <http://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/UCM453398.pdf>
- National Veterinary Assay Laboratory Forestry and Fisheries. (2012). *A Report on the Japanese Veterinary Antimicrobial Resistance Monitoring System 2008 to 2011*. Available at: [http://www.maff.go.jp/nval/tyosa\\_kenkyu/taiseiki/pdf/jvarm2008\\_2011.pdf](http://www.maff.go.jp/nval/tyosa_kenkyu/taiseiki/pdf/jvarm2008_2011.pdf)
- Rhodes, G., Huys, G., Swings, J., McGann, P., Hiney, M., Smith, P., et al. (2000). Distribution of oxytetracycline resistance plasmids between aeromonads in hospital and aquaculture environments: implication of Tn1721 in dissemination of the tetracycline resistance determinant *tet A*. *Appl. Environ. Microbiol.* 66, 3883–3890. doi: 10.1128/AEM.66.9.3883-3890.2000
- Rodriguez-Mozaz, S., Chamorro, S., Marti, E., Huerta, B., Gros, M., Sánchez-Melsió, A., et al. (2015). Occurrence of antibiotics and antibiotic resistance genes in hospital and urban wastewaters and their impact on the receiving river. *Water Res.* 69, 234–242. doi: 10.1016/j.watres.2014.11.021
- Sanchez-Cespedes, J., Figueras, M. J., Aspiroz, C., Aldea, M. J., Toledo, M., Alperi, A., et al. (2009). Development of imipenem resistance in an *Aeromonas veronii* biovar *sobria* clinical isolate recovered from a patient with cholangitis. *J. Med. Microbiol.* 58, 451–455. doi: 10.1099/jmm.0.47804-0
- Schwaiger, K., Holzel, C., and Bauer, J. (2010). Resistance gene patterns of tetracycline resistant *Escherichia coli* of human and porcine origin. *Vet. Microbiol.* 142, 329–336. doi: 10.1016/j.vetmic.2009.09.066
- Segura, P. A., Takada, H., Correa, J. A., El Saadi, K., Koike, T., Onwona-Agyeman, S., et al. (2015). Global occurrence of anti-infectives in contaminated surface waters: impact of income inequality between countries. *Environ. Int.* 80, 89–97. doi: 10.1016/j.envint.2015.04.001

## AUTHOR CONTRIBUTIONS

Study conception and design: MU, CB, SS, HT, YT. Acquisition of data: MU, CT, AF, TO, KS, HT. Analysis and interpretation of data: MU, CT, SS, HT, YT. Drafting of manuscript: MU, SS, HT, YT.

## FUNDING

This work was supported by MEXT KAKENHI Grant Number 25257402.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.00710>

- Shimizu, A., Takada, H., Koike, T., Takeshita, A., Saha, M., Rinawati, et al. (2013). Ubiquitous occurrence of sulfonamides in tropical Asian waters. *Sci. Total Environ.* 45, 108–115. doi: 10.1016/j.scitotenv.2013.02.027
- Smith, P., Hiney, M. P., and Samuelsen, O. B. (1994). Bacterial resistance to antimicrobial agents used in fish farming: a critical evaluation of method and meaning. *Ann. Rev. Fish Dis.* 4, 273–313. doi: 10.1016/0959-8030(94)90032-9
- Suzuki, S., and Hoa, P. T. (2012). Distribution of quinolones, sulfonamides, tetracyclines in aquatic environment and antibiotic resistance in indochina. *Front. Microbiol.* 3:67. doi: 10.3389/fmicb.2012.00067
- Suzuki, S., Ogo, M., Koike, T., Takada, H., and Newman, B. (2015). Sulfonamide and tetracycline resistance genes in total- and culturable-bacterial assemblages in South African aquatic environments. *Front. Microbiol.* 6:796. doi: 10.3389/fmicb.2015.00796
- Tamminen, M., Karkman, A., Lohmus, A., Muziasari, W. I., Takasu, H., Wada, S., et al. (2011). Tetracycline resistance genes persist at aquaculture farms in the absence of selection pressure. *Environ. Sci. Technol.* 45, 386–391. doi: 10.1021/es102725n
- Yano, Y., Hamano, K., Tsutsui, I., Aue-Umneoy, D., Ban, M., and Satomi, M. (2015). Occurrence, molecular characterization, and antimicrobial susceptibility of *Aeromonas* spp. in marine species of shrimps cultured at inland low salinity ponds. *Food Microbiol.* 47, 21–27. doi: 10.1016/j.fm.2014.11.003

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Usui, Tagaki, Fukuda, Okubo, Boonla, Suzuki, Seki, Takada and Tamura. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.